

Exercise and Frailty

Anne Shumway-Cook, PT, PhD
Associate Professor
Department of Rehabilitation Medicine
And
Health Promotion Research Center
University of Washington
Seattle, WA

Objectives

To discuss characteristics of frail older adults and consider the impact of these characteristics on physical activity and exercise prescription.

I. Frailty

A. Defining frailty – many different definitions

- Older adults with a high number of chronic conditions and illnesses
- Older adults who require assistance in activities of daily living
- Frailty characterized by a decrease in reserve capacity and increased risk for disability (Buchner and Wagner, 1992)

B. Pathogenesis of frailty

- Low-grade physiologic loss resulting from a sedentary lifestyle
- Rapid loss due to acute insults (illness, injuries, major life events) that result in periods of limited activity and bedrest.
- To some extent, frailty is preventable

C. Project Homestretch

- Joint project between Department of Rehabilitation Medicine, University of Washington and Aging and Disability Services
- Developed a home based exercise program for frail older adults with Type 2 diabetes
- Comprehensive evaluation and instruction in a home based exercise program targeting: strength, flexibility, balance and gait, aerobic conditioning
- Two year project, 24 ADS clients and 24 healthy older adults

D. Characteristics of frail older adults compared to healthy older adults

- Reduced physical activity and exercise
- Increase in the number and severity of comorbidities
- Increase in the number of medications
- Medically fragile, unstable hypertension, blood sugars
- Required assistance for activities of daily living
- Increased falls and fall risk
- Poor Outcomes Expectation – not all believe in the importance of exercise for their health and function

- Poor Self Efficacy – don't believe that they can exercise safely
 - Fear of falling
 - Fear of injury
- Motivation - fewer “reasons” to exercise – many homebound with limited participation in community activities
- Reduced social support for exercise (e.g. often alone, limited friends available to exercise with)

II. Exercise Programs for Frail Older Adults

1. Components of exercise program are the same

- Strength
- Flexibility
- Posture/Balance
- CV Endurance

2. Begin more slowly, progress more slowly

- No weights
- Short duration
- Low level of exertion
- Modify program in response to change in medical status

3. Modify exercises themselves

- More done in sitting
- Decreased weights
- Emphasis on balance and gait training to reduce falls

4. Instruction on how to exercise safely

- Correct form to prevent injury
- Set up environment for safety
- Prevent falls

5. Monitor response to exercise carefully

- Perceived exertion
- HR
- BP
- Blood sugar

6. Importance of education

- Motivation
- Outcomes Expectation
- Self-Efficacy

IV. Conclusions

- Exercise is essential to all older adults, but especially to frail older adults
- Basic components of an exercise program are unchanged

- Modify exercises
- Monitor response to exercise
- Motivate and educate

References

Bassey EJ, Fiatarone MA, O'Neill EF, et al. Leg extension power and functional performance in very old men and women. Clin Sci (Colch).1992;52:321-327.

Buchner DM, Wagner EH. Preventing frail health. Clin Geriatr Med 1992 Feb;8(1):1-17.

Bortz WM. A conceptual framework of frailty: A review. J of Gerontology: Medical Sciences 2002;57A(5):M283-M288.

Fiatarone -Singh MA. Exercise comes of age: rationale and recommendations for a geriatric exercise prescription. J of Gerontology: Medical Sciences 2002;57A(5):M262-M282.